



## ***In vitro* rumen fermentation characteristics of substrate mixtures with soybean meal partially replaced by microbially fermented yellow wine lees**

K. Y. Yao, F. F. Gu & J. X. Liu

To cite this article: K. Y. Yao, F. F. Gu & J. X. Liu (2020) *In vitro* rumen fermentation

## *In vitro* rumen fermentation characteristics of substrate mixtures with soybean meal partially replaced by microbially fermented yellow wine lees

K. Y. Yao, F. F. Gu and J. X. Liu

### ABSTRACT

This study was conducted to evaluate the effects of replacing soybean meal (SBM) with unfermented and microbially fermented yellow wine lees (YWL) mix on the *in vitro* rumen fermentation characteristics of substrate mixtures. Both types consisting of YWL at 400 g/kg were included in the mixtures at different ratios (1:0, 1:1, 1:2, 1:5 and 0:1, w/w) to replace SBM. Microbial fermentation of YWL did not have a negative impact on gas value ( $p > .05$ ), increased microbial protein (MCP,  $p < .01$ ) and *in vitro* crude protein (CP) digestibility ( $p < .01$ ), and improved the efficiency of nitrogen utilisation ( $p < .01$ ). The ratio of YWL replacing SBM had linear and quadratic effects on gas production (GP,  $p < .01$ ), the rate of GP ( $p < .01$ ), MCP ( $p < .01$ ), total volatile fatty acids (VFAs,  $p < .01$ ), *in vitro* CP digestibility and efficiency of nitrogen utilisation ( $p < .01$ ), with the optimal ratio at 1:1. The *in vitro* digestibilities of dry matter and organic were slightly ( $p < .01$ ) reduced with the increasing ratio of YWL. Besides, positive associative effects were observed on GP parameters, VFA and MCP at some replacement ratios. Considering the *in vitro* rumen characteristics and their associative effects, the optimal ratio at which to replace SBM with microbially fermented YWL was indicated to be 1:1.

### HIGHLIGHTS

- The *in vitro* rumen fermentation characteristics performed better with microbially fermented than unfermented yellow wine lees (YWL) replacing soybean meal (SBM).
- The positive associative effects were observed on gas production parameters, volatile fatty acid and microbial protein at replacing ratio of 1:1.
- The optimal ratio at which to replace SBM with microbially fermented YWL was 1:1.

### ARTICLE HISTORY

### KEY WORDS

*in vitro*

## Introduction

Soybean meal (SBM) is the most important plant protein source in animal industry (Boguhn et al. 2008). In recent years, alternative protein feeds have attracted much interest all over the world, with growing consumer demand for dairy products because of the insufficient supply and high price of SBM. Many studies have been conducted on the utilisation of unconventional feed resources, such as palm meal, distiller's grains and sesame cake, in dairy production (Boguhn et al. 2008). In China, million tons of distiller's grains, by-products of the wine industry, are not well treated, but still contain quite a lot of unused protein, starch, fat and other ingredients which may have feeding value (Schingoethe et al. 2009).

Yellow wine is a traditional Chinese alcoholic beverage made from rice, sorghum or wheat (Hu et al.

2014). The yield of yellow rice wine amounted to 16 billion litres in 2015, generating nearly 1 million tons of yellow wine lees (YWL) as a co-product (Wang 2016). They contain a high content of crude protein (CP, 315–413 g/kg of DM) and the price is one third of that of SBM (Hu et al. 2014). However, YWL has an unbalanced amino acid (AA) profile and is hard to preserve because of its high moisture content (Cheng and Qian 2007). A previous study showed that through microbial pre-treatment, the CP and peptide contents, AA profile and the *in vitro* rumen digestibility of YWL was also improved (Yao et al. 2018), indicating that fermented YWL has potential as protein feed in ruminant rations.

Dairy rations with microbially fermented feed change the rumen fermentation, including increasing rumen microorganisms, enhancing nutrient flow and

### CONTACT



improving the digestibility of feeds (Piamphon et al. 2017). In addition, microbially fermented feed can greatly improve the health and production performance of goats and beef cattle (Syngai et al. 2016). The inclusion of microbially fermented feed in a concentrate mixture at a level of 50% significantly improved the nutrient digestibility of feed and the average daily gain of goats and enhanced their disease resistance (Zhang 2017). A diet with yeast cultures could improve milk production and digestibility of CP and DM in lactating cows (Meller et al. 2019). *Bacillus subtilis* probiotics can increase the degradation rate of feed by *in vitro* fermentation, promote the growth of rumen microorganisms and increase the production of microbial protein (MCP) and volatile fatty acids (VFAs) in the



statistical analyses, the model included substrates, ratio and their interaction as fixed effects. The linear and quadratic effects of the ratio were analysed using orthogonal polynomial contrasts. The cubic effects of the ratio were not examined for inexplicability in biology. Probability values of  $p < .05$  were used to define statistical significance, and values of  $p \leq .10$  and  $p \geq .05$  were regarded as statistical trends.

## **Results and discussion**

### ***Gas production parameters***

The cumulative GP was similar between the UM and FM, but differed among the ratios of UM or FM replacing SBM (Table 2). The GP data of both substrates showed a similar quadratic pattern, i.e. initial increase and then decline with decreasing SBM ratio ( $p < .01$ ). No differences were observed between the two substrates in the rapid degradation fraction of the GP ( $p > .05$ ), while linear and quadratic

2018), contributing to the higher ruminal ammonia-N level in FM than in UM at each replacement. The MCP concentration was higher in the FM than in the UM ( $p < .01$ ). A linear decline in MCP values was observed with increasing levels of UM or FM ( $p < .01$ ), and quadratic effects of the replacement ratio were observed in both substrates ( $p < .01$ ). Ruminal MCP provides 40–80% of the protein needs for ruminants (Uyeno et al.



- Elghandour MMY, Chagoyán JCV, Salem AZM, Kholif AE, Castañeda JSM, Camacho LM, Cerrillo-Soto MA. 2014. Effects of *Saccharomyces cerevisiae* at direct addition or pre-incubation on *in vitro* gas production kinetics and degradability of four fibrous feeds. *Ital J Anim Sci.* 13: 295–301.
- Elghandour MMY, Chagoyán JCV, Salem AZM, Kholif AE, Castañeda JSM, Cipriano MM, Camacho LM, Márque O. 2017. *In vitro* gas and methane production of two mixed rations influenced by three different cultures of *Saccharomyces cerevisiae*. *J Appl Anim Res.* 45:389–395.
- Hu WL, Wu YM, Liu JX, Guo YQ, Ye JA. 2005. Tea saponins affect *in vitro* fermentation and methanogenesis in faunated and defaunated rumen fluid. *J Zhejiang Univ Sci.* 6: 787–792.
- Hu Y, Pan L, Dun Y, Peng N, Liang Y, Zhao S. 2014. Conversion of yellow wine lees into high-protein yeast culture by solid-state fermentation. *Biotechnol Biotechnol Equip.* 28:843–849.
- Ismail SA, Abdel-Fattah AM, Emran MA, A a HH, El-Gamal MS, Hashem AM. 2018. Effect of partial substitution of ration's soybean meal by biologically treated feathers on rumen fermentation characteristics (*in vitro*). *Pak J Biol Sci.* 21:110–118.
- Jia P, Cui K, Ma T, Wan F, Wang W, Yang D, Wang Y, Guo B, Zhao L, Diao Q. 2018. Influence of dietary supplementation with *Bacillus licheniformis* and *Saccharomyces cerevisiae* as alternatives to monensin on growth performance, antioxidant, immunity, ruminal fermentation and microbial diversity of fattening lambs. *Sci Rep.* 8:16712.
- Khampa S, Chaowarat P, Koatdoke U, Singhalert R, Wanapat M. 2009. Manipulation of rumen ecology by malate and yeast in native cattle. *Pak J Nutr.* 8:1048–1051.
- Lawes Agricultural Trust. 1991. GenStat-Release 3.1. VSN International Ltd., Rothamstead. The United Kingdom.
- Lin Y. 2009. The combination effect of distillers' grains, corn and soybean meal and their effects on rumen fermentation characteristics [Doctoral Dissertation]. Ya'an, China: Sichuan Agricultural University.
- Liu JX, Susenbeth A, Südekum K-H. 2002. *In vitro* gas production measurements to evaluate interactions between untreated and chemically treated rice straws, grass hay, and mulberry leaves. *J Anim Sci.* 80:517–524.
- Makkar HPS, Sharma OP, Dawra RK, Negi SS. 1982. Simple determination of microbial protein in rumen liquor. *J Dairy Sci.* 65:2170–2173.
- Mauricio RM, Mould FL, Dhanoa MS, Owen E, Channa KS, Theodorou MK. 1999. A semi-automated *in vitro* gas production technique for ruminant feedstuff evaluation. *Anim Feed Sci Technol.* 79:321–330.
- Meller RA, Wenner BA, Ashworth J, Gehman AM, Lakrit J, Firkins JL. 2019. Potential roles of nitrate and live yeast culture in suppressing methane emission and influencing ruminal fermentation, digestibility, and milk production in lactating Jersey cows. *J Dairy Sci.* 102:6144–6156.
- Piamphon N, Wachirapak C, Bannasan K, Pornsopin P, Sotawong P, Gunun P. 2017. Influence of *Aspergillus niger* or *Saccharomyces cerevisiae*-fermented Napier Grass (*Pennisetum purpureum*) mixed with fresh cassava root on blood parameters and nutrient digestibility in growing beef cattle. *Pak J Nutr.* 16:776–781.
- Rodionow A, Flessa H, Ka ansky O, Guggenberger G. 2006. Organic matter composition and potential trace gas production of permafrost soils in the forest tundra in northern Siberia. *Geoderma.* 135:49–62.
- Schingoethe DJ, Kalscheur KF, Hippen AR, Garcia AD. 2009. Invited review: the use of distillers products in dairy cattle diets. *J Dairy Sci.* 92:5802–5813.
- Syngai GG, Gopi R, Bharali R, Dey S, Lakshmanan GA, Ahmed G. 2016. Probiotics – the versatile functional food ingredients. *J Food Sci Technol.* 53:921–933.
- Togtokhbayar N, Cerrillo MA, Rodrigue GB, Elghandour MMY, Salem AZM, Urangkaich C, Jigjidpurev S, Odongo NE, Kholif AE. 2015. Effect of exogenous xylanase on rumen *in vitro* gas production and degradability of wheat straw. *Anim Sci J.* 86:765–771.
- Uyeno Y, Akiyama K, Hasunuma T, Yamamoto H, Yokokawa H, Yamaguchi T, Kawashima K, Itoh M, Kushibiki S, Hirako M. 2017. Effects of supplementing an active dry yeast product on rumen microbial community composition and on subsequent rumen fermentation of lactating cows in the mid-to-late lactation period. *Anim Sci J.* 88:119–124.
- Valentin SF, Williams PEV, Forbes JM, Sauvart D. 1999. Comparison of the *in vitro* gas production technique and the nylon bag degradability technique to measure short- and long-term processes of degradation of maize silage in dairy cows. *Anim Feed Sci Technol.* 78:81–99.
- Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci.* 74: 3583–3597.
- Wang YC. 2016. China wine work report – 2015. *J Wine Tech.* 5:17–29.
- Yang HJ, Xie CY. 2010. Assessment of fibrolytic activities of 18 commercial enzyme products and their abilities to degrade the cell wall fraction of corn stalks in *in vitro* enzymatic and ruminal batch cultures. *Anim Feed Sci Technol.* 159:110–121.
- Yao KY, Zhang TZ, Wang HF, Liu JX. 2018. Upgrading of by-product from beverage industry through solid-state fermentation with *Candida utilis* and *Bacillus subtilis*. *Lett Appl Microbiol.* 67:557–563.
- Zhang L, Chung J, Jiang Q, Sun R, Zhang J, Zhong Y, Ren N. 2017. Characteristics of rumen microorganisms involved in anaerobic degradation of cellulose at various pH values. *RSC Adv.* 7:40303–40310.
- Zhang Z. 2017. Effect of active yeast and fermented feed on rumen fermentation and nutrient digestibility [Doctoral Dissertation]. Huhhot, China: Inner Mongolia Agricultural University.