



# Natural Product Research

Natural Product Research - International Journal of Natural Product Research

Original Article | **Artepillin C, is it a good marker for quality control of Brazilian green propolis?**

Cui-ping Zhang, Xiao-ge Shen, Jia-wei Chen, Xia-sen Jiang, Kai Wang & Fu-liang Hu (2017)

Artepillin C, is it a good marker for quality control of Brazilian green propolis?, *Natural Product Research*, 31:20, 2441-2444, DOI: [10.1080/14786419.2017.1303697](https://doi.org/10.1080/14786419.2017.1303697)

Online to date: <http://dx.doi.org/10.1080/14786419.2017.1303697>

View supplementary material [↗](#)

Published online: 16 Mar 2017.

Submit your article to this journal [↗](#)

Article views: 58

View related articles [↗](#)

View Crossmark data [↗](#)

Full Terms & Conditions of access and use can be found at <http://www.tandfonline.com/action/journalInformation?journalCode=gnpl20>

SHORT COMMUNICATION

## Artepillin C, is it a good marker for quality control of Brazilian green propolis?

Cui-ping Zhang<sup>a</sup>, Xiao-ge Shen<sup>a</sup>, Jia-wei Chen<sup>a</sup>, Xia-sen Jiang<sup>a</sup>, Kai Wang<sup>b</sup> and Fu-liang Hu<sup>a</sup>

<sup>a</sup>College of Animal Sciences, Zhejiang University, Hangzhou, China; <sup>b</sup>Institute of Apicultural Research, Chinese Academy of Agricultural Sciences (CAAS), Beijing, China

### ABSTRACT

As commercialisation of Brazilian green propolis is going on, quality evaluation and authenticity are important. The result demonstrated that artepillin C found by far in Brazilian green propolis by HPLC-ESI-MS/MS analysis, while a small interferent may be mistaken as artepillin C in some propolis from China. A new HPLC quality control method as artepillin C for marker was developed, which is the primary assessment criteria for quality control of Brazilian green propolis.

### 1. Introduction

Brazilian green propolis, its plant origin is *Baccharis dracunculifolia* DC (Kumazawa et al. 2003), has been widely used in health-care, cosmetic and medicine products especially in Japan and China since 1990s. Contrary to poplar-type propolis, Brazilian green propolis has relatively low concentration of flavonoids while phenolic acids especially isoprenyl phenylpropanoids, caffeoylquinic acid and diterpenes are much richer (Salatino (a)0.5.5 (n ofee)0.5 (s)0.5 (

It is generally accepted that artepillin C is the characteristic chemical composition (Park et al. 2004) and has become an important factor in evaluating the quality of Brazilian green propolis. However, Wang (2010) identified artepillin C in Chinese propolis from Jilin province using ESI-MS/MS. Wu et al. (2013) also detected artepillin C by HPLC in Chinese propolis from Shandong, Beijing, Jiangsu and Xijiang provinces. It is necessary to know with certainty whether artepillin C only exists in Brazilian green propolis?

This study is aimed to verify if artepillin C also exist in other propolis samples and develop a simple and selective method for evaluation of artepillin C. Samples of propolis of different geographic origins were investigated. Furthermore, we compared the content of artepillin C concentrations among different Brazilian green propolis. These results will provide scientific theoretical basis for the quality control of Brazilian green propolis.

## 2 Results and discussion

### 2.1. The determination of artepillin C in propolis

The artepillin C was analysed under HPLC conditions from poplar-type propolis, birch-type propolis, eucalyptus-type and macaranga-type propolis (Figure S2). No interfering peaks were found with retention times similar to artepillin C except propolis samples from Jilin, China. There is a small peak near where artepillin C appeared. ESI-MS/MS was carried out in both positive and negative mode. The fingerprint peak in Brazilian green propolis showed very similar ESI-MS2 spectra as the artepillin C. Major negative ion markers are those of  $m/z$  299.8, 255.7 and 200.5, and major positive ion markers are those of  $m/z$  301.7, 283.7, 245.6, 277.5, 177.4 and 69.2 (Figures S3a, S3b). Although the fragment with  $[m/z]$  301.7 in positive ion is a common ion product for artepillin C and the unknown compounds from Chinese propolis and the other major ion markers from artepillin C were not observed in Chinese propolis (Figure S3c). Another experiment revealed that the fragment with  $[m/z]$  301.7 in positive ion was absent in Chinese propolis by Agilent 1200 QTOF LC/MS 6510 (Figure S4).

Propolis fraction is highly complex, misidentification is very possible. This is the case of two recent papers based on HPLC (Wu et al. 2013) and ESI-MS/MS (Wang 2010). HPLC identification of artepillin C is inadequate because it can't exclude the interfering compounds. In fact, in ESI-MS/MS the unknown compound only gives the ion product with  $[m/z]$  300.4, 283.9 (Wang 2010), it also can't exclude the interfering compounds without more information.

### 2.2. The quantitative method of artepillin C by HPLC

A rapid method based on HPLC with UV detection is presented for the quantitative determination of artepillin C in propolis. The calibration curve of artepillin C was obtained by plotting peak areas vs. six known concentrations (0.05–0.5 mg/mL). The representative linear equation obtained was:  $y = 9178.5x - 36.208$  ( $r^2 = 0.9993$ ), which showed good linear relationships between the peak areas and the concentrations. The LOD and LOQ were 0.12 and 0.4  $\mu\text{g/mL}$ , respectively. The obtained values for both LOD and LOQ were low, which indicated that the method was capable of detecting and quantifying trace amounts of artepillin C in samples. The RSD of inter- and intra-batch precisions were 0.35 and 0.79%, respectively; which showed good precision of the method. The repeatability of HPLC analysis for hydro-alcoholic extract was exemplified by an RSD of 0.078%. The accuracy was studied through

recovery tests and obtained in the ranges of 98.46–99.20%. These results indicated that the developed method was validated and applicable for sample analysis.

Artepillin C were analysed in 56 Brazilian green propolis samples, it can be detected in all samples varying from 0.84 to 8.23%, and the average value is 3.09% (Table S1); which is in accordance with the results reported by Park et al. (2004) but lower than the values obtained in Matsuda's study (2008). The significant difference of artepillin C content among different batches might be account of geographic origins, seasonality, and storage condition and so on (Simões-Ambrosio et al. 2010; Bueno-Silva et al. 2016; Talla et al. 2016).

### 3. Conclusion

Propolis samples from different geographical origins were analysed, artepillin C existed by far in Brazilian green propolis. A small interferent may be mistaken as artepillin C in some propolis from China. A HPLC method was developed for quality control over quantity of artepillin C in Brazilian green propolis.

### Acknowledgements

We thank Wu Xiaodan and Zhu Yaer from Center of Analysis & Measurement of Zhejiang University for technical support and assistance.

### Disclosure statement

No potential conflict of interest was reported by the authors.

### Funding

This work was supported by the grant from the National Natural Science Foundation of China under [grant number 31402146] and Modern Agro-industry Technology Research System from the Ministry of Agriculture of China under [grant number CARS-45].

### References

- Bueno-Silva B, Marsola A, Ikegaki M, Alencar SM, Rosalen PL. 2016. The effect of seasons on Brazilian red propolis and its botanical source: chemical composition and antibacterial activity. *Nat Prod Res.* 31:1318–1324.
- Kumazawa S, Yoneda M, Shibata I, Kanaeda J, Hamasaka T, Nakayama T. 2003. Direct evidence for the plant origin of Brazilian propolis by the observation of honeybee behavior and phytochemical analysis. *Chem Pharm Bull.* 51:740–742.
- Matsuda AH, de Almeida-Muradian LB. 2008. Validated method for the quantification of artepillin-C in Brazilian propolis. *Phytochem Anal.* 19:179–183.
- Park YK, Paredes-Guzman JF, Aguiar CL, Alencar S, Fujiwara FY. 2004. Chemical constituents in *Baccharis dracunculifolia* as the main botanical origin of Southeastern Brazilian propolis. *J Agric Food Chem.* 52:1100–1103.
- Salatino A, Teixeira ÉW, Negri G. 2005. Origin and chemical variation of Brazilian propolis. *Evid-based Compl Alt.* 2:33–38.
- Simões-Ambrosio LMC, Gregório LE, Sousa J.P.B., Figueiredo-Rinhel ASG, Azzolini AECS, Bastos JK, Lucisano-Valim YM. 2010. The role of seasonality on the inhibitory effect of Brazilian green propolis on the oxidative metabolism of neutrophils. *Fitoterapia.* 81:1102–1108.

- Talla E, Tamfu AN, Gade IS, Yanda L, Mbafor JT, Laurent S, Elst LV, Popova M, Bankova V. **2016**. New mono-ether of glycerol and triterpenes with DPPH radical scavenging activity from Cameroonian propolis. *Nat Prod Res.* 31:1379–1389.
- Wang W. **2010**. Research on quality evaluation of propolis from different geographic origins. Changchun: Changchun University of Chinese Medicine.
- Wu JQ, GaoWN Wei JY, Pu LL, Jiao CY, Guo CJ. **2013**. Comparative study on the effective components in propolis from different areas. *Food Nut in China.* 19:62–65.