

Effects of extraction methods on the antioxidant activities of polysaccharides obtained from *Flammulina velutipes*



Zuofa Zhang^{a,1}, Guoying Lv^{a,1}, Weiqiang He^b, Liangen Shi^c, Huijuan Pan^a, Leifa Fan^{a,*}

^a Institute of Horticulture, Zhejiang Academy of Agricultural Science, Hangzhou 310021, China

^b Institute of Biology and Environment, Jiaxing Vocational and Technical College, Jiaxing 314036, China

^c College of Animal Science, Zhejiang University, Hangzhou 310029, China

ARTICLE INFO

Article history:

Received 8 May 2013

Received in revised form 25 June 2013

Accepted 22 July 2013

Keywords:

Flammulina velutipes

Antioxidant activity

Polysaccharide

Extraction method

ABSTRACT

Four polysaccharides (CFP, UFP, MFP and EFP) were extracted from *Flammulina velutipes* using hot water, ultrasonic, microwave or enzymatic methods optimized by orthogonal test. Preliminary structural characterizations were conducted using physicochemical properties. Polysaccharides extracted by all four methods showed similar physicochemical characteristics and FT-IR spectra. However, SEM images of tissues of *F. velutipes* were significantly different. EFP demonstrated better antioxidant activities against hydroxyl radical as well as improved metal chelating activity. UFP showed higher DPPH scavenging activity, but CFP exhibited higher antioxidant activity in reducing power. Hence, these polysaccharides can be used as natural antioxidants in functional foods or medicine. Further experiments on the biological activities of these four polysaccharides are currently in progress.

1. Introduction

Antioxidant status in humans reflects the dynamic balance between the antioxidant defense and prooxidant conditions. When the mechanism of antioxidant protection becomes unbalanced by factors such as aging, impaired physiological functioning may occur, resulting in diseases and accelerated aging (Tiwari, 2004). It is widely accepted that appropriate supplementation with exogenous antioxidant may help reduce reactive oxygen species (ROS)-induced oxidant damage. Due to concern about the side-effects of some synthetic antioxidants, there is increasing interest in replacing synthetic antioxidants with natural antioxidants in the food, pharmaceutical and cosmetic industries (Chen, You, Huang, Yu, & Chen, 2010; Tannin-Spitz, Bergman, Van-Moppes, Grossman, & Arad, 2005).

Many reports have indicated that fungal polysaccharides have strong antioxidant activities in general and should be explored as novel potential antioxidants (Song, Zhang, Zhang, & Wang, 2010; Wang et al., 2010). The antioxidant activity of polysaccharides mainly depends on several structural parameters including sugar composition, molecular weight, type of glycosidic bond in the main

chain, and the degree of sulfuric acid esterification (Lu, Wang, Hu, Huang, & Wang, 2008).

Flammulina velutipes is one of the most popular edible fungi in China and Japan because of its high nutritional value and attractive taste. It has been reported that both the fruiting bodies and the fungal mycelia of *F. velutipes* contain bioactive polysaccharides with beneficial immunomodulatory, anti-tumor, and biological activity on hepatocytes, as well as antioxidant activity (Leung, Fung, & Choy, 1997).

In this study, polysaccharides from *F. velutipes* were extracted using hot water, ultrasonic, microwave, and enzyme-based methods. Preliminary structural characterization of the four polysaccharides was then conducted via physicochemical property, Fourier transform infrared (FTIR) spectroscopy, and scanning electron microscopy analyses. Finally, the antioxidant activities of the extracted polysaccharides were determined against DPPH and hydroxyl radicals, in addition to assaying their metal chelating activity and reducing power. The main aim of this study is to investigate the application value of polysaccharides from *F. velutipes*.

2. Materials and Methods

2.1. Materials

The fruiting body of *F. velutipes* was obtained from the farm at the Zhejiang Academy of Agricultural Science. The mushroom were ground to fine powder, then defatted with petroleum ether and pretreated twice with 80% ethanol prior to experiments.

* Corresponding author. Tel.: +86 571 86404319; fax: +86 571 86404319.

E-mail address: fanleifa2009@126.com (L. Fan).

¹ These authors contributed equally to this manuscript and worked as co-first authors.

1,1-Diphenyl-2- picrylhydrazxyl (DPPH), 3-(2-pyridyl)-5, 6-diphenyl-1, 2,4-triazine-4,4-disulfonic acid monosodium salt (ferrozine), butyl hydroxy anisole (BHA), trifluoroacetic acid (TFA), monosaccharide standards (α -galactose, α -arabinose, β -fucose, β -rhamnose, α -mannose, α -xylose, α -glucose) and EDTA were purchased from Sigma (St. Louis, MO, USA). Cellulase (160 U/mg) was supplied by Huzhou Lilai Chemical Reagent Co. Ltd. All other chemicals and solvents were of the highest commercial grade and obtained from Huadong Chemical Reagent Co. Ltd. (Hangzhou, China).

2.2. Extraction methods

2.2.1. Conventional solvent extraction (CSE) of polysaccharides from *F. velutipes*

Approximately 2.0 g of ground powder was mixed with distilled water in a round-bottom flask. The flask was placed into a HH-6 water bath (Guohua Wiring Company, Shanghai, China) and connected to the condenser. Extractions were performed under the conditions specified in [Table 1a](#). The suspension was centrifuged (4000 \times g, 15 min) and the insoluble part of the plant material was treated again as mentioned above. The supernatant was incorporated

Table 1aAnalysis of L9 (3)³ test results of conventional solvent extraction (CSE).

| No. | A, extraction temperature (°C) | B, extraction time (min) | C, ratio of water to raw material | Extraction yield (%) |
|----------------|--------------------------------|--------------------------|-----------------------------------|----------------------|
| 1 | 75 | 60 | 20 | 6.98 |
| 2 | 75 | 90 | 30 | 7.21 |
| 3 | 75 | 120 | 40 | 7.95 |
| 4 | 85 | 60 | 30 | 7.46 |
| 5 | 85 | 90 | 40 | 9.37 |
| 6 | 85 | 120 | 20 | 8.24 |
| 7 | 95 | 60 | 40 | 9.68 |
| 8 | 95 | 90 | 20 | 8.65 |
| 9 | 95 | 120 | 30 | 8.96 |
| K ₁ | 7.38 | 8.04 | 7.96 | |
| K ₂ | 8.36 | 8.41 | 7.88 | |
| K ₃ | 9.09 | 8.38 | 9.00 | |
| R | 1.71 | 0.34 | 1.12 | |

R refers to the result of extreme analysis.

Extraction yield (%)=(polysaccharides weight/mushroom powder weight) × 100.

Table 1bAnalysis of L9 (3)³ test results of ultrasound-assisted extraction (UAE).

| No | A, extraction temperature (°C) | B, extraction time (min) | C, ratio of water to raw material | D, ultrasonic power (W) | Extraction yield (%) |
|----------------|--------------------------------|--------------------------|-----------------------------------|-------------------------|----------------------|
| 1 | 30 | 30 | 20 | 100 | 8.76 |
| 2 | 40 | 30 | 30 | 200 | 10.58 |
| 3 | 50 | 30 | 40 | 300 | 11.46 |
| 4 | 30 | 40 | 40 | 200 | 10.23 |
| 5 | 40 | 40 | 20 | 300 | 10.35 |
| 6 | 50 | 40 | 30 | 100 | 9.56 |
| 7 | 30 | 50 | 30 | 300 | 10.98 |
| 8 | 40 | 50 | 40 | 100 | 9.77 |
| 9 | 50 | 50 | 20 | 200 | 9.36 |
| K ₁ | 9.99 | 10.27 | 9.49 | 9.36 | |
| K ₂ | 10.23 | 10.05 | 10.37 | 10.06 | |
| K ₃ | 10.12 | 10.04 | 10.48 | 10.93 | |
| R | 0.24 | 0.23 | 0.99 | 1.33 | |

R refers to the result of extreme analysis.

Extraction yield (%)=(polysaccharides weight/mushroom powder weight) × 100.

at 562 nm. The percentage inhibition of ferrozine–Fe²⁺ complex formation was determined using the following formula:

$$\text{Chelating activity (\%)} = \left(\frac{A_0 - A_1}{A_0} \right) \times 100,$$

where A_0 is the absorbance of the control, and A_1 is the absorbance of the samples. The control contained FeCl₂ and ferrozine complex formation molecules. EDTA was used as positive control.

2.4.3. Reducing power

The reductive potential of all samples was determined by the method described by Oyaizu (1986). Different concentrations of test samples (0.31–5.00 mg/mL) were mixed with phosphate buffer

(2.5 mL, 0.2 M, pH 6.6) and potassium ferricyanide [K₃Fe(CN)₆] (2.5 mL, 1%). The mixture was incubated at 50 °C for 20 min. Trichloroacetic acid (2.5 mL, 10%) was added to the mixture, which was then subjected to centrifugation (10 min, 1000 × g). The upper layer of solution (2.5 mL) was removed and mixed with distilled water (2.5 mL) and FeCl₃ (0.5 mL, 0.1%), and the absorbance was measured at 700 nm.

2.4.4. Hydroxyl radical scavenging assay

Hydroxyl radical-scavenging activity was measured according to Smirnoff and Cumbes's work (1989). For this, 0.5 mL FeSO₄ (1.5 mM) was mixed with 0.35 mL H₂O₂ (6 mM), 0.15 mL sodium salicylate (20 mM) and 1.0 mL sample (0.31–5.00 mg/mL), then

Table

Table 1dAnalysis of L9 (3)⁴ test results of enzymatic method for extraction (EAE).

| No | A, extraction time (min) | B, extraction temperature (°C) | C, ratio of water to raw material | D, cellulase concentration (wt.% of mushroom powder) | Extraction yield (%) |
|----------------|--------------------------|--------------------------------|-----------------------------------|--|----------------------|
| 1 | 60 | 40 | 20 | 0.5% | 7.26 |
| 2 | 60 | 50 | 30 | 1.0% | 6.80 |
| 3 | 60 | 60 | 40 | 1.5% | 7.94 |
| 4 | 120 | 40 | 30 | 1.5% | 7.01 |
| 5 | 120 | 50 | 40 | 0.5% | 7.45 |
| 6 | 120 | 60 | 20 | 1.0% | 8.15 |
| 7 | 180 | 40 | 40 | 1.0% | 7.71 |
| 8 | 180 | 50 | 20 | 1.5% | 8.07 |
| 9 | 180 | 60 | 30 | 0.5% | 8.90 |
| K ₁ | 7.33 | 7.33 | 7.83 | 7.87 | |
| K ₂ | 7.54 | 7.44 | 7.57 | 7.95 | |
| K ₃ | 8.23 | 8.33 | 7.70 | 7.67 | |
| R | 0.9 | 1.0 | 0.26 | 0.28 | |

R refers to the result of extreme analysis.

Extraction yield (%) = (polysaccharides weight/mushroom powder weight) × 100.

incubated at 37 °C for 1 h. The absorbance of the hydroxylated salicylate complex was measured at 562 nm. BHA was used as the positive control. The antioxidant activity was calculated with the following equation:

$$\text{Scavenging effect (\%)} = \left(\frac{A_0 - A_1}{A_0} \right) \times 100$$

where A_0 was the absorbance of the solvent control, and A_1 is the absorbance of the test (sample or BHA).

2.5. Statistical analysis

All the experiments were carried out in triplicate and data were expressed as mean ± standard deviation. Statistical analysis was performed with one-way analysis of variance (ANOVA). Statistical significance of differences between the groups was assessed by Student's *t*-test. All the calculations were performed in the SAS 9.0. A level of $P < 0.05$ was taken as statistically significant.

3. Results and discussion

3.1. Optimization of the extraction conditions of the four methods

An orthogonal test was used to optimize the four extraction methods (CSE, UAE, MAE and EAE). **Table 1** shows the polysaccharide yield obtained under the experimental conditions evaluated. With regard to CSE, extraction temperature had the highest *R* value (1.71) and exerted the greatest effect on the polysaccharide yield. The effects of the variables on extraction yields followed the order: extraction temperature > ratio of water to raw material > extraction time. With regard to UAE, ultrasonic power exhibited the greatest effect on the extraction yield, as evidenced by the highest *R* value (1.33), the effects of the variables on extraction yields followed the order: ultrasonic power > ratio of water to raw material > extraction temperature > extraction time. With regard to MAE, microwave power had the highest *R* value (2.03) and exerted the greatest effect on the polysaccharide yield, the effects of the variables on extraction yields followed the order: microwave power > extraction temperature > ratio of water to raw material > extraction time. With regard to EAE, extraction temperature and time exhibited higher *R* values (1.0 and 0.9, respectively). The effects of the variables on extraction yields followed the order: extraction temperature > extraction time > cellulase concentration > ratio of water to raw material.

As shown in **Table 1**, the optimum conditions for each process were: CSE at ratio of water to raw material of 40:1, an extraction temperature of 95 °C, and an extraction time of 90 min; UAE at ratio

Table 2

Comparison of the yields and basic physicochemical characteristics of polysaccharides of the four methods.

| Method | CSE | MAE | UAE | EAE |
|---------------------------------------|----------------|-------|-------|-------|
| Extraction temperature (°C) | 95 | 110 | 30 | 60 |
| Extraction power (W) | – | 500 | 300 | – |
| Extraction time (min) | 90 | 10 | 40 | 180 |
| Ratio of water to raw material (mL/g) | 40 | 30 | 40 | 20 |
| Yield (%) | 9.68 | 12.67 | 11.46 | 9.02 |
| Polysaccharide content (%) | 58.63 | 52.58 | 60.10 | 64.31 |
| Protein content (%) | 1.64 | 3.86 | 1.36 | 0.96 |
| Sugar components (mol%) | | | | |
| L-Fucose | 25.34 | 18.49 | 15.31 | 6.45 |
| D-Mannose | 18.20 | 20.54 | 24.35 | 19.81 |
| D-Glucose | 21.35 | 31.46 | 37.86 | 37.89 |
| D-Galactose | 35.11 | 29.35 | 22.48 | 35.85 |
| Phenol-sulfuric acid test | + ^a | + | + | + |
| α-Naphthol reaction | + | + | + | + |
| Iodination reaction | – ^b | – | – | – |
| Fehling's test | – | – | – | – |
| Carbazole reaction | – | – | – | – |
| FeCl ₃ reaction | – | – | – | – |
| Coomassie brilliant blue reaction | + | + | + | + |

^a Positive.^b Negative.

of water to raw material of 40:1, an extraction temperature of 40 °C, extraction power of 300 W and an extraction time of 30 min; MAE at ratio of water to raw material of 30:1, an extraction temperature of 110 °C, extraction power of 500 W, and an extraction time of 10 min; EAE at ratio of water to raw material of 20:1, an extraction temperature of 60 °C, an extraction time of 180 min, and an enzyme concentration of 1.0% (wt.% of mushroom powder).

3.2. Physicochemical property analysis

Table 2 shows the efficiencies of CSE, MAE, UAE and EAE for extraction of polysaccharide from the fruit body of *F. velutipes*. MAE and UAE exhibited higher efficiencies with high yields and short extraction times, compared to CSE and EAE. Considerable pressure was built up inside the materials during the MAE process (Kratchanova, Pavlova, & Panchev, 2004). This high pressure drastically changed the physical properties of the cell wall, breaking down the structure of cells and improving the capillary-porous structure of treated tissues. High-intensity shock waves generate intense pressures, shear forces and temperature gradient due to the bubble of cavitation; indeed, this is responsible for the majority of physical, chemical and mechanical effects of ultrasound within a material, which can rupture plant cell walls and accelerate mass transfer into the solution, increasing the extraction efficiency (Chen




Fig. 1. Infrared spectra of the four polysaccharides from *Flammulina velutipes*.

et al., 2012). Enzymatic hydrolysis disrupts cell walls, accelerates the migration of target compounds from the material to the surroundings and increases the contact area; these effects enhance the polysaccharide extraction efficiency (Li, Zhang, Han, & Row, 2012). The heated solvent used in CSE diffused slowly through the cell walls, and dissolved and carried away the target compound. Thus, the microstructure of tissues was slightly damaged.

Neutral sugar content was highest in EFP (64.31%) then followed by UFP (60.10%), CFP (58.63%) and MFP (52.58%). Each extract contained protein and its concentration decreased in the following order: MFP > CFP > UFP > EFP. Analysis of sugar composition was performed on the four polysaccharides by GC indicated that they all consist of L-fucose, D-mannose, D-glucose and D-galactose. As shown in Table 2, the results from the phenol-sulfuric acid, α -naphthol, iodination, Fehling's, carbazole, FeCl₃, and Coomassie brilliant blue tests were similar for all four polysaccharides. These similarities indicated that the four extractions were polysaccharides that contained some proteins, but did not contain starch, reducing sugar, uronic acid, or polyphenols. In conclusion, the basic physicochemical properties of the four polysaccharides were similar.

3.3. FTIR analysis

FTIR spectroscopy is typically used for the qualitative measurement of organic functional groups, especially O—H, N—H, and C=O. Fig. 1 shows the FTIR spectra of the four polysaccharides from *F. velutipes*. There were stretching vibrations of O—H and saturated C—H at 3300–3500 and 2927–2930 cm^{−1}, respectively. Two amide bands, one at around 1640 cm^{−1} for Amide I (for C=O) indicated that the four polysaccharides had conjugated proteins. The group of bands extending from 1485 cm^{−1} to 1350 cm^{−1} were assigned to —CH (O—CH₂) flexural vibrations. The absorption band at 1000–1200 cm^{−1} suggested that the four polysaccharides contained pyranose monomers in their structures. The bands in the range of 350–600 cm^{−1} were assigned to skeletal modes of pyranose rings (Yang & Zhang, 2009).

3.4. SEM analysis

The extraction efficiency was related to physical changes in the cell wall of the plant tissue. The microstructures of *F. velutipes* tissue after extraction were observed by SEM image. The results in Fig. 2 showed that different extraction methods produced different physical changes. The cell wall of the *F. velutipes* tissue (Fig. 2A and D)

was almost intact after treated by CSE and ESE. However, cell walls treated by UAE and MAE (Fig. 2B and C) were explosively damaged. It was in agreement with previous investigations (Balachandran, Kentish, Mawson, & Ashokkumar, 2006; Ying, Han, & Li, 2011). The SEM analysis provided strong evidence of the high polysaccharide extraction efficiencies of the UAE and MAE processes.

3.5. Antioxidant activity

3.5.1. DPPH scavenging effect

The DPPH radical is a stable free radical and can accept an electron or hydrogen radical to become a stable diamagnetic molecule. It has been widely accepted as a tool for estimating the free-radical scavenging activities of antioxidants (Hu, Lu, Huang, & Ming, 2004). This assay can accommodate many samples in a short period and is sensitive enough to detect active ingredients at low concentrations (Sanchez-Moreno, 2002).

In this study, DPPH was used to determine the proton-scavenging activity of the four polysaccharides (CFP, UFP, MFP and EFP) extracted from *F. velutipes*. The dose-response curves for the four tested samples are shown in Fig. 3A. The four polysaccharides demonstrated clear scavenging activity at all tested concentrations, and their activity correlated well with concentration, increasing up to 5.0 mg/mL. UFP and MFP showed the higher activity. At 2.50 mg/mL, the scavenging activities of CFP, UFP, MFP, EFP and BHA were 58.50%, 66.75%, 64.34%, 57.93% and 99.82%, respectively. Ultrasonic and microwave treatments may induce the degradation of polysaccharide, which was supposed that the chemical groups in degrade sample have more chance to contact with radical because of the better water-soluble and more surface area (Zhang, Wang, Mo, & Qi, 2013). The effect of antioxidants on DPPH scavenging was thought to be due to their hydrogen-donating abilities. In this study, the tested samples showed scavenging activity on DPPH, which may be attributable to a strong hydrogen-donating ability.

3.5.2. Reducing power

A direct correlation between the antioxidant and reducing capacity has been reported (Amarowicz, Pege, Rahimi-Moghaddam, Barl, & Weil, 2004). Reducing properties are generally associated with the presence of reductones, which can donate a hydrogen atom and exert antioxidant action by breaking the free-radical chain (Gordon, 1990). Reductones are also reported to react with certain precursors of peroxide, preventing peroxide formation (Duh, Du, & Yen, 1999). Thus, the reducing capacity of a compound or extract may be a significant indicator of its potential antioxidant activity (Fan, Li, Deng, & Ai, 2012). The reducing powers for all samples are shown in Fig. 3B, the absorbance was 0.058–0.174 for CFP, 0.068–0.158 for UFP, 0.053–0.126 for MFP and 0.054–0.099 for EFP. However, the reducing power of the four tested samples was much weaker when compared with the positive control, BHA. Reducing properties are generally associated with the presence of electron-donating groups or hydrogen atoms. These data on reducing power of the tested polysaccharides indicated that it could play a role in the antioxidant observed.

3.5.3. Metal chelating assay

Iron-chelating may produce important antioxidant effects by retarding metal-catalyzed oxidation (Kehrer, 2000). Effective Fe²⁺ chelators may also afford protection against oxidative damage by removing Fe²⁺ that may otherwise participate in HO[·]-generating Fenton-type reactions. Minimizing Fe²⁺ may also protect against oxidative damage by inhibiting the production of reactive oxygen species and lipid peroxidation. The Fe²⁺-chelating capacity of the tested samples was determined by measuring iron–ferrozine complexes (Fig. 3C). The Fe²⁺ chelating effects of all






Fig. 2. Scanning electron micrographs of *Flammulina velutipes* after extraction ($3000\times$): (A) treated by CSE (conventional solvent extraction); (B) treated by UAE (ultrasound-assisted extraction); (C) treated by MAE (microwave-assisted extraction) and (D) treated by EAE (enzymatic method for extraction).

polysaccharides were concentration-dependent. The Fe^{2+} chelating effect of EFP was the highest than the other polysaccharides, though all of them had lower activity than EDTA in the concentration range from 0.31 to 5.00 mg/mL. The chelating capability of the

polysaccharides decreased as follows: EFP > CFP > MFP > UFP. Various polysaccharides exhibited remarkable binding capacity for Fe^{2+} , further demonstrating they had stronger antioxidant capability. Therefore, we hypothesize that the ferrous ions chelating



effects of the tested

- Macdonald, J., Galley, H. F., & Webster, N. R. (2003). Oxidative stress and gene expression in sepsis. *British Journal of Anaesthesia*, 90, 221–232.
- Oyaizu, M. (1986). Studies on product of browning reaction prepared from glucose amine. *Japanese Journal of Nutrition*, 44, 307–315.
- Sanchez-Moreno, C. (2002). Methods used to evaluate the free radical scavenging activity in foods and biological systems. *Food Science and Technology International*, 8, 121–137.
- Sheng, J., Yu, F., Xin, Z., Zhao, L., Zhu, X., & Hu, Q. (2007). Preparation, identification and their antitumor activities in vitro of polysaccharides from *Chlorella pyrenoidosa*. *Food Chemistry*, 105, 533–539.
- Shimada, K., Fujikawa, K., Yahara, K., & Nakamura, T. (1992). Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *Journal of Agricultural and Food Chemistry*, 50, 840–845.
- Smirnoff, N., & Cumbes, O. J. (1989). Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry*, 28, 1057–1060.
- Song, H. F., Zhang, Q. B., Zhang, Z. S., & Wang, J. (2010). In vitro antioxidant activity of polysaccharides extracted from *Bryopsis plumose*. *Carbohydrate Polymers*, 80, 1057–1061.
- Song, M., Li, Q. X., & He, C. L. (2009). Identification and analgesic effect of *Angelica Sinensis* extract. *Journal of Xiangning University*, 23, 194–196.
- Spector, J. (1978). Refinement of the comassie blue method of protein qualification. A simple and linear spectrophotometric assay of 0.5–50 µg protein. *Analytical Biochemistry*, 86, 142–146.
- Tannin-Spitz, T., Bergman, M., Van-Moppes, D., Grossman, S., & Arad, M. S.