

Genetic characterization of H9N2 avian influenza virus in plateau pikas in the Qinghai Lake region of China

Yan Yan¹ · Jin-Yan Gu² · Zhuang-Chuan Yuan¹ · Xin-Yu Chen¹ · Zeng-Kui Li³ ·
Jing Lei² · Bo-li Hu² · Li-Ping Yan² · Gang Xing² · Min Liao¹ · Ji-Yong Zhou^{1,4}

Received: 7 June 2016 / Accepted: 10 November 2016 / Published online: 18 December 2016
© Springer-Verlag Wien 2016

Abstract Qinghai Lake is a major migratory-bird breeding site that has experienced several highly pathogenic avian influenza virus (AIV) epizootics. Plateau pikas (*Ochotona curzoniae*) have previously been implicated in the ecology of avian influenza virus in this region. We first isolated an H9N2 AIV (A/Pika/Menyuan/01/2008) from plateau pikas between November 2008 and October 2009. Sequence analysis showed that the A/Pika/Menyuan/01/2008 AIV was closely related to the H9N2 AIV strain (A/Turkey/Wisconsin/1/1966). Our findings suggested that plateau pikas may contribute to AIV epidemiology in the Qinghai Lake region.

Keywords Qinghai Lake · Plateau pika · Avian influenza virus · H9N2

Introduction

H9N2 avian influenza virus (AIV) was first isolated in Guangdong Province of China between November 1992 and May 1994 [1]. Following the initial emergence of this virus in China, widespread transmission of H9N2 AIV was observed in chickens, but vaccination campaigns helped to control the virus in poultry. Currently, H9N2 AIV is endemic in multiple avian species in different regions. Based on the hemagglutinin (HA) gene, H9N2 virvirusocanTL9.96259

Materials and methods

Sample collection and virus identification

To identify AIV infections in wild pikas at Qinghai Lake between November 2008 and October 2009, tissue samples were collected from plateau pikas in Haibei, Huangnan, Hainan, Haixi, Guoluo and Yushu Prefecture of Qinghai Province (Fig. 1). Tissue samples were stored at -80°C prior to inoculation into 10-day-old specific-pathogen-free (SPF) embryonated chicken eggs and propagation (3 passages). The allantoic fluids with positive hemagglutination activity were harvested and stored at -80°C . Subtyping of the positive samples was done by RT-PCR [5, 6].

Genome sequence analysis

The genomic sequences of the virus isolates were determined as described previously [6]. Briefly, total RNA was extracted from the allantoic fluid using TRIzol[®] LS Reagent (Invitrogen, Carlsbad, CA, USA), and RT-PCR was then performed using a OneStep RT-PCR Kit (QIAGEN, Hilden, Germany) with

primers specific for the H9N2 virus HA gene HA-F 5', CTCAGGGAGCAAAAGCAGGGG-3'; HA-R 5', GGCATGGCCCAGAATAGGAAG-3') and the NA gene NA-F 5', ATGAATCCAAATCAGAAG-3'; NA-R 5' CTACTTGTCAATGGTGAATG-3'). The PCR products were cloned into the pMD18-T vector and then sequenced. Multiple sequence alignments were made using the Clustal W program of the BioEdit software package (version 5.0.9). Phylogenetic analysis was carried out using MEGA (version 4).

Nucleotide sequence accession number

The nucleotide sequence of the H9N2 virus in this study has been deposited in the GenBank database (accession number shown in Table 1).

Results

Only one virus was isolated from 817 tissue samples collected from 138 pikas. The virus was isolated from the kidney tissue of one pika in Menyuan County of Qinghai



Fig. 1 Plateau pika sampling sites in Qinghai Province, China, between 2008 and 2009. Round red symbols indicate sites of pika sample collection. The pink point indicates the site of the isolated H9N2 virus. QL, Qilian; WL, Wulan; DLH, Delingha; GEM,

Ge'ermu; ZD, Zhiduo; YS, Yushu; JZ, Jiuzhi; MD, Maduo; GH, Gonghe; MY, Menyuan; CD, Chengduo; JZH, Jianzha; QML, Qumalai (color figure online)

Table 1 Genetic similarity of A/Pika/Menyuan/01/2008 (H9N2) to the most closely related H9N2 viruses

Gene	Virus with the highest nucleotide sequence identity	Source	Identity	GenBank accession no.
HA	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99.4%	CY014663
	A/chicken/Heilongjiang/35/2000(H9N2)	Avian	98.9%	DQ064366
NA	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99%	CY087826
PB2	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99.8%	CY014670
PB1	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99.5%	CY014669
PA	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99.6%	CY014668
NP	A/chicken/Heilongjiang/35/2000(H9N2)	Avian	99.6%	DQ064447
M	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99.9%	CY014664
NS	A/turkey/Wisconsin/1/1966(H9N2)	Avian	99%	CY087828

Province in December 2008. Subtype analysis showed that the isolated virus belonged to the H9N2 subtype of avian influenza virus, and it was named A/Pika/Menyuan/01/2008 (H9N2, Pk/MY/1/08). After three passages in chicken eggs, the HA titer of the Pk/MY/01/08 was 2^3 .

Sequence analysis showed that the Pk/MY/1/08 virus contained the motif PAVSSR/GL at the cleavage site between HA1 and HA2, which is a characteristic of low-pathogenic avian influenza virus (LPAI). Phylogenetic analysis demonstrated that the HA of Pk/MY/1/08 shared 99.4% and 98.9% nucleotide sequence identity with that of A/Turkey/Wisconsin/1/1966 (TY/WI/66) and A/Chicken/Heilongjiang/35/00, respectively (Fig. 2A). Similar findings were also confirmed for the NA gene, which shared 92.6%–99% nucleotide sequence identity with viruses of the North American TY/Ca/189/66-like lineage (Fig. 2B). There were no amino acid deletions in the stem of the NA protein of the Pk/MY/1/08 virus.

Phylogenetic analysis indicated that all of the internal genes of Pk/MY/1/08 had the genetic characteristics of members of the TY/Ca/189/66-like lineage (data not shown). The internal genes of Pk/MY/1/08 showed high nucleic sequence similarity to those of the Ty/Ca/189/66-like strains. The polymerase genes (PB2, PB1 and PA), M genes, and NS genes shared 99.5%–99.8%, 99.9%, and 99% nucleotide sequence identity, respectively, with those of the A/Turkey/Wisconsin/1/1966, whereas the NP gene shared 99.6% identity with that of the A/chicken/Heilongjiang/35/2000. Residues 26–34 of the M2 protein correspond to the LVIAASIIG motif found in ion channels, and no mutation was found at position 31, which is associated with the resistance phenotype of influenza A virus. The presence of 627E and 701D in PB2 indicated that Pk/MY/1/08 is a strain with low virulence that has not acquired mutations for adaptation to the mammalian host. Like those of the pika H5N1 viruses isolated in 2005, the NS gene of Pk/MY/1/08 belongs to the Y439-like lineage.

Discussion

Qinghai Lake is an important breeding site for migratory birds in the central Asian-India flyway. Therefore, infected wild birds may transmit AIVs to other birds or mammals in the Qinghai Lake region. Plateau pikas live in close proximity to wild birds in this ecosystem. Following the first outbreak of H5N1 HPAIV in migratory birds at Qinghai Lake in 2005, we detected HPAI H5N1 viruses in pikas around Qinghai Lake [6]. This finding raises concern that plateau pikas may play a role in the transmission of AIVs at Qinghai Lake and provide a potential opportunity for AIVs to adapt to mammals. Notably, other groups have demonstrated that wild lagomorphs are susceptible to multiple subtypes of influenza A virus [8, 9] and that some of them have the potential to transmit these viruses back to waterfowl [10].

Researchers from the United States and China have shown that wild birds migrate from Qinghai Lake to Mongolia, Eastern Asia, and the Bay of Bengal [11]. Qinghai Lake, which is the largest salt lake in China, is an important breeding site for migratory birds on the Central Asia-India migratory route. A previous study showed evidence of HPAI H5N1 virus in plateau pikas and suggested that this lagomorph may have the potential to transmit AIVs to domestic mammals and humans [6]. Therefore, we speculate that pikas may serve as a potential spillover host of AIV in the Qinghai Lake region and may have the potential to transmit influenza A viruses to other species. Notably, others have shown that cottontail rabbits (*Sylvilagus* sp.) were able to transmit avian AIV to mallards that shared their living space [10].

Five H9N2 influenza viruses were isolated from the environment at a live poultry market in the Qinghai Lake region in 2012 [12]. In our study, we isolated the Pk/MY/1/08 strain of AIV, which has high sequence similarity to members of the North American lineage A/Turkey/Wisconsin/1/1966. However, the reason for this similarity is unclear. A similar result was found in a previous study, in




Fig. 2 Phylogenetic trees of the HA and NA genes of H9N2 avian influenza viruses. The phylogenetic trees show that HA (A) and NA (B) of the *A/Pika/Menyuan/01/2008* strain belong to the Ty/Ca/1/66-like lineage. The phylogenetic trees were generated in MEGA 4 using

the neighbor-joining (NJ) algorithm with 1,000 bootstrap replicates, and the NJ bootstrap values (>70%) for each node are shown in each tree. The virus characterized in this study is indicated by a red circle (color figure online)

which the *A/CK/Heilongjiang/35/00* (H9N2) strain, which contains HA and NP genes of the North American lineage, was isolated [12]. Interestingly, the Pk/MY/1/08 virus was isolated from kidney tissue. Therefore, pikas with asymptomatic AIV infections may provide potential opportunities

for the adaptation of avian influenza viruses to mammals and may also be involved in the epidemiology of the viruses in the Qinghai Lake region. Additional studies are needed to evaluate the potential role of plateau pikas in the transmission of AIVs in this region.

Acknowledgements This study was supported by the National Key Technology R&D Program of China (Grant No. 2016YFC1201604, 2015BAD12B00) and the Wildlife Department of the State Forestry Administration.

Compliance with ethical standards

Conflict of interest All authors declare that they have no conflict of interest.

References

1. Sun YP, Liu JH (2015) H9N2 influenza virus in China: a cause of concern. *Protein Cell* 6(1):18–25
2. Wang B, Chen QJ, Chen Z (2012) Complete genome sequence of an H9N2 avian influenza virus isolated from Egret in Lake Dongting wetland. *J Virol* 86(21):11939
3. Zhang K, Zhang ZW, Yu ZJ, Li L, Cheng KH, Wang TC, Huang G, Yang ST, Zhao YK, Feng N, Fu J, Qin CA, Gao YW, Xia XZ (2013) Domestic cats and dogs are susceptible to H9N2 avian influenza virus. *Virus Res* 175(1):52–57
4. Liu J, Xiao H, Lei F, Zhu Q, Qin K, Zhang XW, Zhang XL, Zhao D, Wang G, Feng Y, Ma J, Liu W, Wang J, Gao GF (2005) Highly pathogenic H5N1 influenza virus infection in migratory birds. *Science* 309(5738):1206
5. Zhou JY, Shen HG, Chen HX, Tong GZ, Liao M, Yang HC, Liu JX (2006) Characterization of a highly pathogenic H5N1 influenza virus derived from bar-headed geese in China. *J Gen Virol* 87:1823–1833
6. Zhou JY, Sun WB, Wang JH, Guo JQ, Yin W, Wu NP, Li LJ, Yan Y, Liao M, Huang Y, Luo KJ, Jiang XT, Chen HL (2009) Characterization of the H5N1 highly pathogenic avian influenza virus derived from wild Pikas in China. *J Virol* 83(17):8957–8964
7. Yu ZJ, Cheng KH, Sun WY, Xin Y, Cai JS, Ma RL, Zhao QB, Li L, Huang J, Sang XY, Li X, Zhang K, Wang TC, Qin C, Qian J, Gao YW, Xia XZ (2014) Lowly pathogenic avian influenza (H9N2) infection in Plateau pika (*Ochotona curzoniae*), Qinghai Lake, China. *Vet Microbiol* 173(1–2):132–135
8. Root JJ, Shriner SA, Bentler KT, Gidlewski T, Mooers NL, Spraker TR, VanDalen KK, Sullivan HJ, Franklin AB (2014) Shedding of a low pathogenic avian influenza virus in a common synanthropic mammal—the cottontail rabbit. *PLoS One* 9:e102513
9. Root JJ, Bosco-Lauth AM, Bielefeldt-Ohmann H, Bowen RA (2016) Experimental infection of peridomestic mammals with emergent H7N9 (A/Anhui/1/2013) influenza A virus: implications for biosecurity and wet markets. *Virology* 487:242–248
10. Root JJ, Shriner SA, Ellis JW, VanDalen KK, Sullivan HJ, Franklin AB (2015) When fur and feather occur together: inter-class transmission of avian influenza A virus from mammals to birds through common resources. *Sci Rep* 5:12354
11. Prosser D (2012) Wild birds and emerging diseases: modeling avian influenza transmission risk between domestic and wild birds in China. *Dissertations & Theses—Gradworks*
12. Li CJ, Yu KZ, Tian GB, Yu DD, Liu LL, Jing B, Ping JH, Chen HL (2005) Evolution of H9N2 influenza viruses from domestic poultry in Mainland China. *Virology* 340(1):70–83