

# Identification of the Paneth cells in chicken small intestine

L. Wang,<sup>1</sup> J. Li,<sup>1</sup> J. Li Jr., R. X. Li, C. F. Lv, S. Li, Y. L. Mi, and C. Q. Zhang<sup>2</sup>

*Department of Veterinary Medicine, College of Animal Sciences, Zhejiang University, Hangzhou 310058, P.R. China*

**ABSTRACT** The Paneth cells are highly specialized cells in the epithelium of the small intestine of many vertebrate species. These cells reside at the base of crypts of the Lieberkühn and contain abundant secretory granules. Previous studies suggesting the existence of Paneth cells in the chicken (*Gallus gallus*) remained controversial. Here we seek to identify the Paneth cells in the chicken small intestine through morphological examination and specific gene expression. Histological staining and transmission electron microscope confirmed the presence of granulated secretory

cells at the base of the crypts in the chicken small intestine. Western blotting experiment also manifested the expression of lysozyme protein, which is specifically secreted by the Paneth cells in the small intestine. Moreover, lysozyme *c* and lysozyme *g* mRNAs were expressed in the small intestine of chickens at different ages. Lysozyme *c* mRNA, in particular, was located at the base of the small intestinal crypts as displayed by in situ hybridization. Collectively, we provide evidences that the Paneth cells indeed exist in the small intestine of the chicken.

**Key words:** paneth cell, chicken, crypt, lysozyme

2016 Poultry Science 95:1631–1635  
<http://dx.doi.org/10.3382/ps/pew079>

## INTRODUCTION

In mammals, the Paneth cells play important immune defense roles in the intestines, particularly relating to intestinal epithelium renewal and early stages of intestinal inflammation. The Paneth cells are restricted to the crypts of the small intestine, along with goblet cells, enterocytes, tuft cells, and enteroendocrine cells. These cells represent the principal cell types of the epithelium of the small intestine (van Es and Clevers, 2014). The Paneth cells were first described by J. Paneth in 1888 as granulated cytoplasmic cells located at the base of small intestinal crypts, the “crypts of Lieberkühn”. Classical Paneth cells are pyramidal shaped secretory cells with basally situated nuclei, extensive endoplasmic reticulum and Golgi network, and prominent, large apical granules that occupy most of their cytoplasm. The large secretory granules in the Paneth cells are rich in antimicrobial peptides (such as lysozyme,  $\alpha$ -defensins/cryptdins, and secretory phospholipase A2), immune modulators (such as IgA), and trophic factors (such as carboxylic ester hydrolase) (Porter et al., 2002; Elphick and Mahida, 2005; Ouellette, 2010; BeS9190010.51.1453.8161.3(d)0.1smissiETSalzmancs001scnBT-0.000299999Tc10.50010.5215.52025150.172010

**Table 1.** Primers for PCR analysis.

Gene	Accession no.	Primer sequence (5'-3')	Product length (bp)
Lysozyme <i>c</i>	NM.205281	GACGATGTGAGCTGGCAG GGATGTTGCACAGGTTCC	225
Lysozyme <i>g</i>	X61002	CACGCTGGCAAAATACTGAAG TTCCCAACACCAGCATTGTAG	236
Lysozyme <i>g2</i>	XM.416896	CATTCCATCTTTGGTTGC CCACCTTTGAGCTGCTGTTC	246
$\beta$ -actin	NM.205518	ACACCCACACCCCTGTGATGAA TGCTGCTGACACCTTCACCATTC	136

intestine based upon the examination of morphological characteristics and the expression of the marker genes or proteins.

Lysozyme, a product secreted by the Paneth cells, is a glycosidase that specifically hydrolyses peptidoglycan, a major component of the bacterial cell wall (Ghoos and Vantrappen, 1971; Callewaert and Michiels, 2010). Lysozyme is also widely found in the surface fluid of tears, and in breast milk, saliva, gastric and small intestinal secretions, as well as in the granules of macrophages and neutrophils (Erlandsen et al., 1974; Spicer et al., 1977). There are three types of lysozyme that have been identified: the c-type (chicken or conventional type), which is highly expressed in the oviduct, macrophages, and small intestine (Hindenburt et al., 1974; Nile et al., 2004); the g-type (goose type), which is expressed in the bone marrow and the lung (Nakano and Graf, 1991); and the g2-type, which is identified in the small intestine, liver, and kidney (Nile et al., 2004). In the gastrointestinal tract, lysozyme c is expressed in the gastric and pyloric glands, duodenal

sections were washed in succession with  $2 \times \text{SSC}$  (saline sodium citrate),  $0.5 \times \text{SSC}$ , and  $0.2 \times \text{SSC}$ , blocked with 5% BSA in buffered saline for 30 min, and incubated with biotin-labeled mouse anti-Digoxigenin for 60 min at  $37^\circ\text{C}$ . After a brief wash in PBS, the sections were detected using the 3,3'-diaminobenzidine tetrahydrochloride (**DAB**) system according to the manufacture's protocol (MK 10152, Boster Bioengineering Co., Ltd., Wuhan, China) and then counterstained with hematoxylin.

### Western Blot

Protein was extracted from 0.1 g tissues using 1 mL RIPA Lysis Buffer (P0013B, Beyotime) containing 10  $\mu\text{L}$  100 mM PMSF (ST506, Beyotime). Electrophoresis was carried out by 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (**SDS-PAGE**), followed by electrotransfer to a polyvinylidene fluoride membrane (Millipore, Billerica, MA). The membrane was incubated in 5% skimmed milk/Tris-buffered saline and Tween 20 at room temperature for 1 h and subsequently incubated with polyclonal rabbit anti-human lysozyme EC 3.2.1.17 antibody (1:200, A0099, Dako) and monoclonal mouse anti-avian  $\beta$ -actin antibody (1:200, sc-47778, Santa Cruz) at  $4^\circ\text{C}$  overnight. The rabbit anti-human lysozyme antibody was initially checked for the feasibility in chicken studies through the immunohistological staining and Western blot. After washing in Tris-buffered saline and Tween 20, the membranes were exposed to horseradish peroxidase labeled secondary antibody (goat anti-rabbit or goat anti-mouse IgG, dilution 1:1000) for 1 h at  $37^\circ\text{C}$ . The membranes were washed three times and visualized using an electrochemiluminescence (**ECL**) system (170-5056, BIO-RAD Laboratones. Inc., Shanghai, China). The  $\beta$ -actin band was adopted as the internal control. The bands obtained in the Western blot were scanned and analyzed by image analysis software (Gel-Pro Analyzer 4.5, Media Cybernetics). The data were expressed as the IOD of the bands, normalized to the IOD of the corresponding  $\beta$ -actin bands. The result was obtained from three replicates.

### Data Analysis

The data were expressed as the means  $\pm$  SD. The difference of lysozyme expression in the same intestine segment between mouse and chicken was analyzed by Independent-Samples *t* tests with SPSS11.5 (SPSS Inc., Chicago, IL). Level for determination of significance was 0.05.

## RESULTS

### Secretory Granules Exist in Rodlike Cells at the Base of Small Intestinal Crypts

Granules in Paneth cells can be recognized by phloxine-tartrazine staining. The staining result showed




**Figure 1.** Secretory granules were observed in rodlike cells located at the base of chicken intestinal crypts at 6 months. (A) Phloxine-tartrazine staining of the small intestine villus in chicken at 6 months. Scale bar: 10  $\mu\text{m}$ . (B) Phloxine-tartrazine staining of the small intestine crypt in chicken at 6 months. The arrow shows the suspected Paneth cells. Scale bar: 10  $\mu\text{m}$ . (C) TEM image of the villus showing the ultrastructure of the granules in the goblet cells. Scale bar: 5  $\mu\text{m}$ . (D) TEM image of the crypt showing the ultrastructure of the granules in the rodlike cells (indicated in white dotted lines). Scale bar: 5  $\mu\text{m}$ .

that some purplish red granules could only be observed in the small intestinal crypts but not in the villus (Figure 1A, B) of 6-month-old chickens. This result implied that Paneth cells might exist in the small intestinal crypts of the chicken intestine. Furthermore, TEM results showed that a large amount of secretory granules existed in the rodlike cells at the bottom of the chicken's intestinal crypts (Figure 1D). However, the granules in the goblet cells at villus showed a low electron density and are arranged in a specific shape (Figure 1C) while granules in the crypts presented a higher electron density (Figure 1D), these different characteristics indicated that the secretory cells in villus and crypt were of different types. It seemed likely that such cells, located at the base of the chicken's small intestinal crypt and confirmed as containing secretory granules, were Paneth cells.


### Expression of Lysozyme in the Chicken Small Intestine

To investigate the Paneth cells in the chicken small intestine, we detected the expression of lysozyme by Western blot. As shown in Figure 2, lysozyme protein was expressed in the small intestine of chicken at 6 months old and mouse (positive control) without statistical significance. However, no positive band was detected in skin and muscle of chicken (negative control).

RT-PCR analysis of mRNA showed lysozyme *c* and lysozyme *g* mRNA expression in the intestine from 1 wk, 2 wk, 3 wk, 8 wk, and 6-month-old chickens (Figure 3A). Whilst the expression of lysozyme *g2* in the jejunum and ileum remained uniform from 1 wk to



**Figure 2.** Western blot analysis of lysozyme protein in the chicken of 6 months old and mouse (A), and the statistical analysis result of three replicates (B). Duo, duodenum; Jej, jejunum; Ile, ileum; Mus, muscle.  $\beta$ -actin was adopted as the internal control. There was no statistical significance ( $P < 0.05$ ) in the same intestine segment between the mouse and chicken.



**Figure 3.** Expression of chicken lysozyme mRNA. (A) Expression of lysozyme *c* (*lys c*), lysozyme *g* (*lys g*), and lysozyme *g2* (*lys g2*) mRNAs on Due (duodenum), Jej (jejunum) and Ile (ileum) at the age of 1 wk, 2 wk, 3 wk, 8 wk and 6 months. (B) Negative control of in situ hybridization. (C) The lysozyme *c* gene was detected at the base of the crypts of the 6-month-old chicken intestine. (D) Magnified images from C. Scale bar: 50  $\mu$ m for B and C, and 10  $\mu$ m for D.

8 wk, in the duodenum it could only be detected from 1 wk to 3 wk.

To further confirm the cells at the base of small intestinal crypt to be Paneth cells, in situ hybridization analysis was performed on the small intestinal sections of chickens. The result showed that lysozyme *c* mRNA was expressed at the base of small intestinal crypts, at just the location that was presumed to be that of the Paneth cells (Figure 3C, D).

## DISCUSSION

Paneth cells are specialized epithelial cells of the small intestine that are located at the base of the intestinal crypts of many vertebrate species. In Paneth cells, abundant secretory granules occur which con-

tain lysozyme, secretory phospholipase A2, enteric  $\alpha$ -defensins, cryptdin related sequence peptides, and angiogenin 4 (Bevins and Salzman, 2011). Among them, lysozyme is widely considered to be a marker of Paneth cells (Cadwell et al., 2008; Sato et al., 2009; Yin et al., 2014). To date, the presence of Paneth cells in the small intestine of the chicken has remained controversial. In 1974, Paneth cells in the avian intestine were reported based on the light-microscopic methods (Humphrey and Turk, 1974). However, a study by Nile et al (2004) showed that there were no Paneth cells in chicken intestine since the expression of lysozyme *c* gene at different ages could not be detected. In this study, by using phloxine-tartrazine staining and TEM, abundant secretory granules were observed at the rod-like cells which were located at the bottom of small intestinal crypts of the chicken (Figure 1B, D). Unlike the Paneth cells in mice and humans, these secretory cells in the chicken small intestinal crypts were thin and elongated into rod shapes. Species diversity may be the cause of this difference in morphological characteristics in such cells between the chicken and mouse. However, these results did imply the existence of Paneth cells in the chicken small intestine.

Lysozyme is one of the most widely used markers for the Paneth cells based on its specific expression in the intestinal crypts. Our RT-PCR result revealed that lysozyme *c* and lysozyme *g* mRNAs are expressed in the chicken in different small intestine segments at various ages. Additionally, the lysozyme *c* gene was specifically located at the base of intestinal crypts which is the predicted location of Paneth cells as indicated by the preliminary morphological analysis above. Moreover, the location of lysozyme *c* mRNA expression in the chicken small intestine was consistent with that in humans (Erlandsen et al., 1974). Therefore, the Paneth cells were considered to exist in chicken small intestine.

In conclusion, we detected the Paneth cells in chicken small intestinal crypts by phloxine-tartrazine staining and TEM observation. We also detected the expression of lysozyme mRNA and protein expression in the chicken small intestine. Lysozyme *c* gene was also specifically expressed at the base of the intestinal crypts. Based on these results, we assume the presence of the Paneth cells in the chicken small intestine. However, the exact roles of the chicken Paneth cells involved in the innate immunity and maintenance of intestinal homeostasis requires further studies.

## ACKNOWLEDGMENTS

This study was supported by the National Natural Science Foundation of China (Nos. 31502023 and 31272525), the Fundamental Research Funds for the Central Universities (No. 2014QNA6024) and scientific research project of Zhejiang Bureau of Education (No. Y201533358). We are grateful to Weidong Zeng for help

in the experiments and Dr. Chris Wood (Zhejiang University) for English improvement in the manuscript.

## REFERENCES

- Adolph, T. E., M. F. Tomczak, L. Niederreiter, H. J. Ko, J. Bock, E. Martinez-Naves, J. N. Glickman, M. Tschurtschenthaler, J. Hartwig, S. Hosomi, M. B. Flak, J. L. Cusick, K. Kohno, T. Iwawaki, S. Billmann-Born, T. Raine, R. Bharti, R. Lucius, M. N. Kweon, S. J. Marciniak, A. Choi, S. J. Hagen, S. Schreiber, P. Rosenstiel, A. Kaser, and R. S. Blumberg. 2013. Paneth cells as a site of origin for intestinal inflammation. *Nature*. 503:272–276.
- Bevins, C. L., and N. H. Salzman. 2011. Paneth cells, antimicrobial peptides and maintenance of intestinal homeostasis. *Nat. Rev. Microbiol.* 9:356–368.
- Cadwell, K., J. Y. Liu, S. L. Brown, H. Miyoshi, J. Loh, J. K. Lennerz, C. Kishi, W. Kc, J. A. Carrero, S. Hunt, C. D. Stone, E. M. Brunt, R. J. Xavier, B. P. Sleckman, E. Li, N. Mizushima, T. S. Stappenbeck, and H. W. Virgin. 2008. A key role for autophagy and the autophagy gene *Atg16l1* in mouse and human intestinal Paneth cells. *Nature*. 456:259–263.
- Callewaert, L., and C. W. Michiels. 2010. Lysozymes in the animal kingdom. *J. Biosci.* 35:127–160.
- Clevers, H. C., and C. L. Bevins. 2013. Paneth cells: maestros of the small intestinal crypts. *Annu. Rev. Physiol.* 75:289–311.
- Cunliffe, R. N., F. R. Rose, J. Keyte, L. Abberley, W. C. Chan, and Y. R. Mahida. 2001. Human defensin 5 is stored in precursor form in normal Paneth cells and is expressed by some villous epithelial cells and by metaplastic Paneth cells in the colon in inflammatory bowel disease. *Gut*. 48:176–185.
- Elphick, D. A., and Y. R. Mahida. 2005. Paneth cells: their role in innate immunity and inflammatory disease. *Gut*. 54:1802–1809.
- Erlandsen, S. L., J. A. Parsons, and T. D. Taylor. 1974. Ultrastructural immunocytochemical localization of lysozyme in the Paneth cells of man. *J. Histochem. Cytochem.* 22:401–413.
- Ghoos, Y., and G. Vantrappen. 1971. The cytochemical localization of lysozyme in Paneth cell granules. *Histochem. J.* 3:175–178.
- Hindenburg, A., J. Spitznagel, and N. Arnheim. 1974. Isozymes of lysozyme in leukocytes and egg white: evidence for the species-specific control of egg-white lysozyme synthesis. *Proc. Natl. Acad. Sci. U.S.A.* 71:1653–1657.
- Humphrey, C. D., and D. E. Turk. 1974. The ultrastructure of normal chick intestinal epithelium. *Poult. Sci.* 53:990–1000.
- Klockars, M., and S. Reitamo. 1975. Tissue distribution of lysozyme in man. *J. Histochem. Cytochem.* 23:932–940.
- Nakano, T., and T. Graf. 1991. Goose-type lysozyme gene of the chicken: sequence, genomic organization and expression reveals major differences to chicken-type lysozyme gene. *Biochim. Biophys. Acta*. 1090:273–276.
- Nevalainen, T. J., J. M. Gronroos, and M. Kallajoki. 1995. Expression of group II phospholipase A2 in the human gastrointestinal tract. *Lab. Invest.* 72:201–208.
- Nile, C. J., C. L. Townes, G. Michailidis, B. H. Hirst, and J. Hall. 2004. Identification of chicken lysozyme g2 and its expression in the intestine. *Cell. Mol. Life Sci.* 61:2760–2766.
- Ouellette, A. J. 2010. Paneth cells and innate mucosal immunity. *Curr. Opin. Gastroenterol.* 26:547–553.
- Porter, E. M., C. L. Bevins, D. Ghosh, and T. Ganz. 2002. The multifaceted Paneth cell. *Cell. Mol. Life Sci.* 59:156–170.
- Sato, T., J. H. van Es, H. J. Snippert, D. E. Stange, R. G. Vries, M. van den Born, N. Barker, N. F. Shroyer, M. van de Wetering, and H. Clevers. 2011. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature*. 469:415–418.
- Sato, T., R. G. Vries, H. J. Snippert, M. van de Wetering, N. Barker, D. E. Stange, J. H. van Es, A. Abo, P. Kujala, P. J. Peters, and H. Clevers. 2009. Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature*. 459:262–265.
- Spicer, S. S., R. Frayser, G. Virella, and B. J. Hall. 1977. Immunocytochemical localization of lysozymes in respiratory and other tissues. *Lab. Invest.* 36:282–295.
- van Es, J. H., and H. Clevers. 2014. Paneth cells. *Curr. Biol.* 24:R547–548.
- Yin, X., H. F. Farin, J. H. van Es, H. Clevers, R. Langer, and J. M. Karp. 2014. Niche-independent high-purity cultures of Lgr5+ intestinal stem cells and their progeny. *Nat. Methods*. 11:106–112.