

Viral Infection Affects Sucrose Responsiveness and Homing Ability of Forager Honey Bees

Zhiguo Li¹, Yanping Chen², Shaowu Zhang^{1,3}, Shenglu Chen¹, Wenfeng Li¹, Limin Yan¹, Liangen Shi¹, Lyman Wu⁴, Alex Sohr⁴, Songkun Su^{1,5*}

1 College of Animal Sciences, Zhejiang University, Hangzhou, Zhejiang, China, **2** USDA, ARS Bee Research Laboratory, Beltsville, Maryland, United States of America, **3** ARC Centre of Excellence in Vision Science, Research School of Biology, College of Medicine, Biology and Environment, the Australian National University, Canberra, Australia, **4** College of Computer, Math, and Natural Sciences, University of Maryland, College Park, Maryland, United States of America, **5** College of Bee Science, Fujian Agriculture and Forestry University, Fuzhou, Fujian, China

Abstract

Honey bee health is mainly affected by *Varroa destructor*, virus

IAPV were detected in the heads of these honey bees. The homing ability significantly in comparison to the homing ability of uninfected foragers in the heads may enable the virus to disorder foraging roles functions that are responsible for learning, navigation, and orientation in have a lower response threshold to sucrose and lose their way back to the

Citation: Li Z, Chen Y, Zhang S, Chen S, Li W, et al. (2013) Viral Infection Affects Sucrose Responsiveness and Homing Ability of Forager Honey Bees. *PLoS ONE* 8(10): e77354. doi:10.1371/journal.pone.0077354

Editor: Adrian G Dyer, Monash University, Australia

Received: July 27, 2013; **Accepted:** September 1, 2013; **Published:** October 10, 2013

Copyright: © 2013 Li et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by earmarked funds for Modern Agro-industry Technology Research Project (2012C12004) from the Natural Science Foundation Commission of Zhejiang Province (R3080306) to SKS. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: susongkun@zju.edu.cn

Introduction

Honey bees (*Apis mellifera* L.) play a vital role in global food production and economy [1]. However, honey bee colony losses in recent years have had a devastating effect on the agricultural industry and ecosystems that rely on honey bees for pollination. Colony Collapse Disorder (CCD) is a poorly understood phenomenon in which workers abruptly disappear and do not return to the hive but leave behind live queen and brood in the colony. This occurred during the winter of 2006-2007 in the US and colony losses reported in recent years in other parts of the world [2,3], poses a particular threat to apiculture and agriculture worldwide. A metagenomic survey of microflora in CCD hives showed that Israeli acute paralysis virus (IAPV), a virus that had not been previously reported in the US, displayed a strong correlation with CCD. IAPV was detected in 86% of CCD-affected colonies but in less than 5%

in healthy colonies [4]. As a result, the disease mechanisms of IAPV infection in honey bees have been the subject of extensive research since then.

IAPV is a member of the genus *Aparavirus*, a new class of viruses in the *Dicistroviridae* family [5]. IAPV was first reported in 2004 in Israel where IAPV infected honey bees were found with symptoms of shivering wings, paralysis, and then death outside of the hives. The severe bee mortality caused by IAPV led to heavy losses in Israeli apiculture [6]. Since its first detection, IAPV infection has been reported in many other parts of the world [7-10]. A new study showed that *Varroa destructor* [11], an ectoparasite of the honey bee, has been catastrophic for the beekeeping industry and is an effective vector of IAPV [12]. The number of copies of IAPV in honey bees was positively correlated with the density of *Varroa* mites and the time period of *Varroa* feeding. The association of IAPV and *Varroa* mites could cause increased damage to honey

immobilized in small glass vials individually by chilling for
about 3 minutes. Sixty honey bees from each group were
ybe f ½ e e t % e v i % i n d i x t % o n a l l \ \ M o

n k E o E

I
a
m

P
r res
M rb
bee " o @ l f i ĩ x t ĩ n d

honeybees erb eè =
m. % b




Figure 1. The RFID system used in homing experiment. (A) Two RFID readers were placed at the customized tunnel entrance of a nucleus hive with 3 frames. (B) A honey bee with the RFID tag glued to its thorax carrying pollen on its hind legs returns to the hive.

doi: 10.1371/journal.pone.0077354.g001

in the homing experiment. The mic3-TAG passive 13.56 MHz tags stored a unique 64-bit number (approximately 2.4 mg, 2.0 x 1.6 mm) and readers (Microsensus 2k6 HEAD) were bought from Microsensus [32]. The ID numbers of tags were input into software running on a computer by a USB-Pen prior to being glued to the thoraxes of honey bees. A 30 cm long customized tunnel was linked to the entrance of a nucleus hive, and two readers were attached to the top of the tunnel (Figure 1). The software used for storing tag ID numbers and exporting data from the readers was designed by Sebastian Streit ("Beegroup ID2DB" © Beegroup, Sebastian Streit, 2003) [31].

As the methods described above, honey bees were captured from the nucleus hives and were then immobilized on ice before injection. One group of 20 pollen foragers was injected with 1 µl of IAPV with a 1: 200 dilution in PBS and another group of 20 pollen foragers was injected with 1 µl of PBS as a control. Each pollen forager was then equipped with an RFID tag on the thorax, using shellac glue, to identify the honey bee. Subsequently, the honey bees were restored in cages at room temperature (24–26 °C) and fed ad libitum with 50% (wt/wt) sucrose solution and water for an hour. After that, the honey bees were then placed in a black box and transported 500 meters away from the hive, at which point we then released these honey bees. Honey bees that could not take off within five minutes were discarded from the study [32]. The two readers recorded the entering or leaving of pollen foragers each day after the start of the experiment [32]. We did not stop observing returning honey bees until the last honey bee didn't return to the hive for each individual trial. The homing experiment was carried out in three different nucleus colonies and each homing experiment was repeated three times for each nucleus colony.

Data analysis

The percentage of honey bees showing PER was calculated by the number of honey bees showing PER divided by the total number of honey bees used for data collection each day after the treatment. Differences in responsiveness to sucrose solution between honey bees injected with IAPV and the control group was analyzed by the Fisher exact test. Comparison of survival rates between the two groups of honey bees was analyzed by independent samples t-test (SPSS Statistics 13.0, SPSS Inc., Chicago, IL, USA). Homing ability was assessed based on the number of bees not returning to the hive each day after the treatment. The data regarding homing ability was not normally distributed. Therefore, a non-parametrical Kruskal-Wallis test followed by Dunn's multiple comparison test was used to analyze the difference of homing ability between the two groups of honey bees using the Graphpad Prism 5 (GraphPad Software Inc, San Diego, CA, USA). Differences were considered significant at $p < 0.05$ for all statistic tests and all tests were two-tailed.

Results

Responsiveness of IAPV-infected honey bees to sucrose

The honey bees infected with 1 µl of IAPV (1: 500 dilution) containing approximately 44 copies of IAPV showed similar sucrose responsiveness at days 0, 1 and 2 after injection in comparison to the sham-injected group of honey bees (Figure 2A). However, significant differences in responsiveness to the low sucrose concentrations between the two groups were found at day 3 after injection (Fisher exact test, $p < 0.05$). These honey bees were found to be more responsive to low sucrose concentrations than that of the sham-injected group of honey




Figure 2. Comparisons between IAPV-injected honey bees and PBS-injected honey bees in their responsiveness to sucrose. Sucrose responsiveness of honey bees injected with IAPV with 1: 500 dilution and PBS-injected honey bees were tested at days 0 (A) and 3 (B) after injection. The number of honeybees tested at day 0 post-injection is 60 for each group; the number of honeybees tested at day 3 for IAPV-injected group is 14 and 29 for the PBS-injected group (*= $p<0.05$, **= $p<0.01$); sucrose responsiveness of honey bees injected with IAPV with 1: 200 dilution and PBS-injected honey bees tested at days 0 (C) and 2 (D) after injection. The number of honey bees tested at day 0 post-injection is 60 for each group; the number of honey bees tested at day 2 for IAPV-injected group is 22 and 30 for the PBS-injected group (*= $p<0.05$, **= $p<0.01$).

doi: 10.1371/journal.pone.0077354.g002

bees, and they showed no difference in responsiveness to the high sucrose water (Figure 2B).

There was no difference regarding sucrose responsiveness between honey bees infected with 1 μ l of IAPV (1: 200 dilution) containing approximately 110 copies of IAPV and honey bees injected with PBS at days 0 and 1 after injection (Figure 2C). However, foragers infected with IAPV exhibited significantly higher responsiveness to the low sucrose water than that of foragers injected with PBS at day 2 after injection (Fisher exact test, $p<0.05$), and they showed no difference in responsiveness to the high sucrose water (Figure 2D).

Survival rates of inoculated honey bees

There was no significant difference between honey bees injected with IAPV and the control group in survival rates (Figure 3). The proportion of live honey bees to dead honey bees was 98.1% and 88.0% for the control group and IAPV-infected honey bees at 24 hours after injection (independent samples t-test, $t=2.025$, $df=3.933$, $p>0.05$), and the proportion at 48 hours after injection was 68.1% and 61.4% for the two groups (independent samples t-test, $t=2.012$, $df=6$, $p>0.05$).

Detection of IAPV in heads of honey bees

The exact number of copies of IAPV in heads of honey bees was shown in Figure 4 by relating the Ct values to the standard curve (Figure S1). There were about 48 copies of IAPV in the




Figure 3. Analysis of survival rates. Comparison of survival rates between honey bees injected with IAPV and honey bees of the control groups at 24, 48 hours after injection. The data are expressed as mean \pm SD of four independent experiments.

doi: 10.1371/journal.pone.0077354.g003

heads of honey bees at day 0 post-injection with IAPV; 6.9×10^6 IAPV copies were detected in the heads of honey bees at day 1 post-injection, and the number of IAPV copies in the heads of honey bees at day 2 post-injection was 1.2×10^7 . IAPV was not detected in the heads of honey bees injected with PBS. The number of copies of IAPV increased dramatically at day 1 post-injection, however, the IAPV load exhibited no obvious increase in heads of honey bees at day 2 post-injection compared to day 1 post-injection.

Impairment of the homing ability of IAPV-infected honey bees

The number of honey bees equipped with tags for control animals and IAPV-infected honey bees was 180, respectively. Only 4 honey bees (1.1%) failed to take off within 5 minutes and these honey bees were excluded from analysis. The homing ability of foragers infected with IAPV was depressed significantly in comparison to foragers injected with PBS (Figure 5). There was no significant difference (Kruskal-Wallis test followed by Dunn's multiple comparison test, $p > 0.05$) between the two groups in the percentage of foragers returning

to the hive at days 0, 1 and 4 post-injection. However, significant differences were found at 2 ($p < 0.01$) and 3 ($p < 0.05$) days post-injection between the two groups. The percentage of returning foragers injected with PBS returning at days 0, 1, 2 and 3 post-injection was 81.0%, 68.9%, 58.7% and 46.0%. For the IAPV injection group, the percentage was 79.6%, 59.2%, 2.3% and 0%, respectively.

At day 2 post-injection, there were around 12 foragers injected with PBS returning to the hive compared with almost no IAPV-injected foragers returning to the hive. About 10 foragers were departing and returning to the hive 3 and 4 days after being injected with PBS, however, there were no foragers departing and returning at day 3 or day 4 after being injected with IAPV. Similar results were obtained from three independent experiments and the data provided clear evidence that a viral infection of IAPV in the heads may make honey bees lose their way back to the hive.

Discussion

Previous studies showed that DWV impaired the sucrose responsiveness of pollen foragers [21]. Our results demonstrated that IAPV also affected the sucrose responsiveness of pollen foragers, causing the honey bees to be more responsive to low concentrations of sugar water as a result of that metabolic stress, compared to honey bees without IAPV infection. IAPV and DWV are both common honey bee viruses and may affect behaviors of honey bees in the same way. However, the underlying disease mechanisms of the virus infections warrants further investigation. Additionally, our study of honey bees infected with different concentrations of viral particles showed similar changing patterns towards different




Figure 5. Effects of IAPV on the homing ability of forager honey bees. The abscissa represents days after foragers were injected with PBS and IAPV respectively. The ordinate represents the percentage of foragers departing and returning to the hive per day. *= $p<0.05$, **= $p<0.01$, n.s.= not significant. Error bars show SEM.

doi: 10.1371/journal.pone.0077354.g005

infested foragers, which was explained as a common response to pathogens responsible for honey bee diseases [29,30]. Bortolotti et al. reported that honey bees treated with sub-lethal doses of imidacloprid did not return to the hive and showed decreased foraging activity compared to that of the control [49]. Henry et al. also reported that the homing ability of forager honey bees was impaired by thiamethoxam intoxication using RFID systems [28]. Here, we provided additional evidence that IAPV infection could impair the homing ability of honey bee foragers as well. Because there was no significant difference between IAPV-infected honey bees and honey bees of the control groups in survival rates after 24 and 48 hours post-injection, we can exclude the possibility that the honey bees that were not returning to the hive in the evening at day 2 post-injection (approximately after 48 hours injection) were primarily due to death caused by mechanism damage. In fact, honey bees that were infected with IAPV at day 2 post-injection departed from the hive in the morning and some honey bees also foraged normally during the daylight hours, however, they did not return to the hive in the evening and they may have gotten lost and died later in the field before returning to the

hive. There were also no dead honey bees found in and outside the hives during the following days. PBS treated honey bees, however, foraged every day, albeit, with the gradual reduction in the number of honey bees returning to the hive during the following days even eighteen days after injection.

A limited number of IAPV copies were detected in the heads of foragers injected with IAPV at day 0 post-injection, and the number of IAPV copies increased rapidly in the heads of foragers on day 1 post-injection. However, the data also show little difference regarding the number of IAPV copies between day 1 post-injection and day 2 post-injection in the heads of foragers injected with IAPV. We concluded that IAPV may interfere with normal nervous system functions in the brains of honey bees and cause foragers to lose their way back to the hives based on the homing behaviors of IAPV-injected foragers and the number of IAPV copies detected in the heads of honey bees. In addition, detection of the viral infection in the heads of honey bees from commercial apiaries was rare, based on our preliminary experiments. Previous studies revealed that the detection of DWV in honey bees' heads is rare and represents an overt DWV infection which is also a significant indicator for

colony loss associated with DWV infection [44,50,51]. Given the fact that there were about 6.9×10^6 and 1.2×10^7 IAPV copies detected in the heads of honey bees at days 1 and day 2 post-injection, respectively, we propose that the high levels of IAPV replication might lead to severe disease in infected honey bees, which lost their navigational abilities and were unable to return to the hives.

An impressive result was found during a homing experiment carried out on a rainy day. There were 14 and 13 foragers for IAPV-injected and PBS-injected control group at day 1 post-injection respectively. All the IAPV-injected foragers that foraged on that day did not return to the hive in the evening. However, only 5 PBS-injected foragers foraged on that day, and 4 of those foragers returned to the hive in the evening. Therefore, 12 PBS-injected foragers foraged outside while the number of IAPV-injected foragers was 0 at day 2 post-injection (a sunny day). Foragers infected with IAPV foraged more actively in adverse weather conditions than the sham-injected foragers did, however, they did not return to the hive. A similar phenomenon was also found in honey bees infected by *Nosema* sp., which was interpreted, in part, as compensation for foraging yield because of their shortened lifespan [30]. Regardless, forager honey bees infected with IAPV at day 2 post-injection showed a trend of foraging earlier in the morning compared to that of the control group. Previous studies showed that foragers with lower response thresholds collect lower concentrations of nectar than those with higher response thresholds [52,53], which can be used to explain why foragers infected with IAPV departed the colony earlier in the day and also foraged on the rainy day.

Our studies clearly showed that both sucrose responsiveness and homing ability of forager honey bees were affected by IAPV which was injected into the hemolymph of honey bees using a 5 μ l microsyringe. The wound and stress caused by the needle might simulate the process of a mite's bite. So, it is conceivable that the stress caused by the needle might have some subtle effects on the honey bees. We can't rule out the possibility that microsyringes used to administer injections might impose additional stress on the honey bees, but it is an effective method to infect the viruses in honey bees [21]. Sucrose responsiveness reflects the division of foraging

labor of honey bees [54] and homing involves spatial memory and navigation of honey bees [26,27]. Viral infection in heads may cause disorders in foraging roles of honey bees, with honey bees foraging abnormally, and may enable the virus to interfere with brain functions that are responsible for navigation, orientation and spatial memory in the honey bees. After foraging, IAPV-infected honey bees initiating homing flight from the foraging site to the hive may lose their way back to the hive due to loss in spatial memory. Our results provided first evidence that viral infection in the heads of honey bees could impair the homing ability of forager honey bees. This study is in line with previous studies that sublethal dosages of insecticides (imidacloprid, thiamethoxam et al.) could affect homing ability and foraging activity of honey bees [28,33,49,55]. Colony losses reported worldwide in recent years can, therefore, be triggered in part by multiple stressors including insecticides and viruses.

Supporting Information

Figure S1. The standard curve for IAPV obtained using SYBR Green qPCR and serial diluted plasmid as template. (TIF)

Acknowledgements

We would like to thank beekeepers for their kindness in providing colonies; Yunjuan Hu, Qin Chen and Ruilu Chen for their assistance in research. We also thank the anonymous referees for many helpful comments, which helped improve this manuscript. ARC Centre of Excellence in Vision Science provided RFID equipment and tags.

Author Contributions

Conceived and designed the experiments: SKS SWZ YPC. Performed the experiments: ZGL WFL LMY. Analyzed the data: SKS YPC ZGL LW SLC. Contributed reagents/materials/analysis tools: SWZ SLC LGS SKS. Wrote the manuscript: ZGL YPC AS SKS.

References

- Gallai N, Salles J-M, Settele J, Vaissière BE (2009) Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecol Econ* 68: 810–821. doi:10.1016/j.ecolecon.2008.06.014.
- VanEngelsdorp D, Underwood R, Caron D, Hayes J Jr (2007) An estimate of managed colony losses in the winter of 2006–2007: A report commissioned by the apiary inspectors of America. *Am Bee J* 147: 599–603.
- Neumann P, Carreck NL (2010) Honey bee colony losses. *J Apic Res* 49: 1–6. doi:10.3896/IBRA.1.49.1.01.
- Cox-Foster DL, Conlan S, Holmes EC, Palacios G, Evans JD et al. (2007) A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* 318: 283–287. doi:10.1126/science.1146498. PubMed: 17823314.
- Chen YP, Nakashima N, Christian P, Bonning BC, Valles SM, Lightner D (2012) Dicistroviridae. In: AM KingMJ AdamsEB CarstensEJ Lefkowitz. *Virus Taxonomy*, 9th. Elsevier Academic Press. pp. 840–845.
- Maori E, Lavi S, Mozes-Koch R, Gantman Y, Peretz Y et al. (2007) Isolation and characterization of Israeli acute paralysis virus, a dicistrovirus affecting honeybees in Israel: evidence for diversity due to intra- and inter-species recombination. *J Gen Virol* 88: 3428–3438. doi: 10.1099/vir.0.83284-0. PubMed: 18024913.
- Blanchard P, Schurr F, Celle O, Cougoule N, Drainudel P et al. (2008) First detection of Israeli acute paralysis virus (IAPV) in France, a dicistrovirus affecting honeybees (*Apis mellifera*). *J Invertebr Pathol* 99: 348–350. doi:10.1016/j.jip.2008.07.006. PubMed: 18703069.
- Yan X, Liu XJ, Chen JH, Zhang Y, Han RC (2009) First detection of Israel acute paralysis virus (IAPV) in *Apis mellifera* (Hymenoptera: Apidae) in China. *Sociobiology* 54: 95–100.
- Chen Y, Evans JD (2007) Historical presence of Israeli acute paralysis virus in the United States. *Am Bee J* 147: 1027–1028.
- Al-Abbadi AA, Hassawi DS, Abu-Mallouh SA, Al-Mazra'awi MS (2010) Novel detection of Israel acute paralysis virus and Kashmir bee virus from honeybees *Apis mellifera* L. (Hymenoptera: Apidae) of Jordan using reverse transcriptase PCR technique. *Appl Entomol Zool* 45: 183–190. doi:10.1303/aez.2010.183.
- Anderson DL, Trueman JW (2000) *Varroa jacobsoni* (Acari: Varroidae) is more than one species. *Exp Appl Acarol* 24: 165–189. doi:10.1023/A: 1006456720416. PubMed: 11108385.

12. Di Prisco G, Pennacchio F, Caprio E, Boncristiani HF Jr, Evans JD et al. (2011) *Varroa destructor* is an effective vector of Israeli acute paralysis virus in the honeybee, *Apis mellifera*. J Gen Virol 92: 151-155. doi:10.1099/vir.0.023853-0. PubMed: 20926637.
13. Maori E, Paldi N, Shafir S, Kalev H, Tsur E et al. (2009) IAPV, a bee-affecting virus associated with Colony Collapse Disorder can be silenced by dsRNA ingestion. Insect Mol Biol 18: 55-60. doi:10.1111/j.1365-2583.2009.00847.x. PubMed: 19196347.
14. Hunter W, Ellis J, Hayes J, Westervelt D, Glick E et al. (2010) Large-scale field application of RNAi technology reducing Israeli acute paralysis virus disease in honey bees (*Apis mellifera*, Hymenoptera: Apidae). PLOS Pathog 6: e1001160.
15. Scheiner R (2004) Responsiveness to sucrose and habituation of the proboscis extension response in honey bees. J Comp Physiol A 190: 727-733. PubMed: 15185117.
16. Bitterman ME, Menzel R, Fietz A, Schäfer S (1983) Classical conditioning of proboscis extension in honeybees (*Apis mellifera*). J Comp Psychol 97: 107-119. doi:10.1037/0735-7036.97.2.107. PubMed: 6872507.
17. Hammer M, Menzel R (1995) Learning and memory in the honeybee. J Neurosci 15: 1617-1630. PubMed: 7891123.
18. Decourtie A, Devillers J, Cluzeau S, Charreton M, Pham-Delègue M-H (2004) Effects of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. Ecotoxicol Environ Saf 57: 410-419. doi:10.1016/j.ecoenv.2003.08.001. PubMed: 15041263.
19. Thany SH, Gauthier M (2005) Nicotine injected into the antennal lobes induces a rapid modulation of sucrose threshold and improves short-term memory in the honeybee *Apis mellifera*. Brain Res 1039: 216-219. doi:10.1016/j.brainres.2005.01.056. PubMed: 15781066.
20. Si A, Zhang S-W, Maleszka R (2005) Effects of caffeine on olfactory